

Waterpipe tobacco smoking

© 2025 National Centre for Smoking Cessation and Training (NCSCT)

Authors: Mohammed Jawad, Andy McEwen, Ann McNeill & Lion Shahab

Editors: Andrew Preston, Melanie McIlvar, Andy McEwen

Date of last review: March 2025

This briefing is based upon an article in the journal *Addiction*¹ and has been prepared by the NCSCT to provide stop smoking practitioners with a summary of the evidence and guidance on the information they can give to their clients.

Executive summary


- Waterpipe tobacco smoking is increasing and the UK has reported a 210% rise in the number of waterpipe cafes nationwide over the last five years.
- Despite a widespread belief amongst people who use waterpipes to smoke tobacco that it is a low risk activity, waterpipe tobacco smoke is known to contain significant levels of toxins including some which are known to be carcinogenic (i.e. causes cancer).
- There is evidence of significant negative health effects from waterpipe tobacco smoking.
- Waterpipe tobacco smoking therefore represents both an individual health issue for clients attending stop smoking services, and a potential public health concern.

Waterpipe tobacco smoking

Waterpipe smoking: history and practice

Waterpipe smoking is a broad term given to a device whereby smoke (usually sourced from a tobacco mixture) is passed through water prior to inhalation.²

Figure 1: Example of a waterpipe with schematic¹



The origins of waterpipe tobacco smoking are thought to date back several centuries, although its exact beginning is uncertain.³

There are a variety of cultural and regional variations in its use, thus it is synonymously known across the world by other terms such as hookah, shisha, narghile, goza or hubble bubble.³ This range of cultural names is partly reflective of the different types of tobacco used.

The most commonly used waterpipe tobacco is called the *Mo'assel* which means 'honeyed' and is made up of about one-third tobacco mixed with two-thirds honey and fruit flavours.⁴ It is separated from heated coal in the waterpipe by pierced aluminium foil.

There are also unflavoured waterpipe tobacco forms known as *Ajami*, *Tumbak*, or *Jurak*, which may have a higher tobacco ratio, and be in direct contact with the heated coal⁴ in waterpipes.

Waterpipe tobacco smoking

The Chinese variant of waterpipe contains very little tobacco, and no coal.⁵ More recently the waterpipe industry has marketed two new variants: firstly, a non-tobacco, herbal variant, claiming to provide a healthier alternative to the *Mo'assel* tobacco⁶, and secondly electronic waterpipe devices attempting to emulate nicotine vapes.⁷

Understanding these differences is important when critiquing the available evidence on the health effects of waterpipe use, as each type of tobacco mixture has a different smoke composition of varying concentrations with slightly different health effects.⁸

Most of the recent literature generally refers to the fruit-flavoured *Mo'assel* type, and it is this type of waterpipe that is discussed in this briefing, unless stated otherwise.

Unlike cigarettes, a waterpipe is commonly smoked intermittently and in groups.^{9–12} It provides a central activity at social gatherings, and is usually smoked for 45–60 minutes¹³, but sometimes longer.¹⁴

The United Kingdom has reported a 210% rise in the number of waterpipe cafes nationwide over the last five years;¹⁵ a regular location for people who smoke. However, smoking is also commonplace in the home^{11,12,16,17} as this is considered cheaper and less time consuming.¹⁸ Waterpipe tobacco use can be associated with other forms of substance abuse, most commonly cigarettes but also recreational drugs,^{9,19,20–23} with reports that narcotics and alcohol can be mixed with the waterpipe tobacco and water respectively to add to its psychoactive effect.^{18,24}

Prevalence and epidemiology

Currently good epidemiological trend data on waterpipe smoking is lacking, in part due to the absence of its inclusion in national health surveys.²⁵

The full picture is further complicated by the fact that waterpipe users do not generally self-identify as people who smoke²⁶ resulting in a tobacco-using population that healthcare professionals are not able to routinely identify.²⁷

Whilst males are more likely to smoke waterpipes than females,²⁸ this gender difference is smaller than that seen for cigarette use, especially in countries where there is a usual gender divide in cigarette use.²⁹ There is no clear trend on socio-economic status predicting waterpipe use, but it may be more prevalent among the more educated and affluent.^{30–32}

Recent longitudinal data suggests an upwards trend which persist across all age groups, but is particularly noticeable in the younger population. In the UK the current prevalence is thought to be 8% amongst school pupils³³ and 8–19% amongst university students.^{25,34,35}

Waterpipe tobacco smoking

Health consequences of waterpipe tobacco smoking

Controlling for the extensive habitual variables of waterpipe smoking such as breathing patterns, size of the waterpipe,³⁶ and materials used,³⁷ has proven difficult when conducting smoke aerosol studies using machine-replicated human behaviour models.³⁸

However, there has been consistency in the production of significant noxious chemicals including tar, carbon monoxide, nitric oxide and various carcinogens.

Attempts have been made to compare these toxicant yields to the amount produced by a single cigarette. Interestingly, despite being marketed as a healthier alternative, it is shown that herbal variants of waterpipe contain significant levels of toxicants, except for nicotine, similar to normal waterpipe *Mo'assel*.

Additionally, one study has shown that waterpipe tobacco delivers substantial amounts of heavy metals including arsenic, beryllium and chromium graded 1 (which is carcinogenic) by the International Agency for Research on Cancer.³⁹

Waterpipe tobacco is known to contain and deliver other carcinogens, e.g. concentrations of radionuclide elements such as uranium, and polonium^{40,41} and tobacco-specific nitrosamines, but in lower amounts than seen in people who smoke cigarettes.⁴²

A key health concern, in the acute setting at least, is the threat of CO poisoning^{43–50} where people generally experience non-specific neurological signs such as light headedness and nausea.¹¹ Waterpipe users usually recall a 'buzz' when smoking, which may reflect the early stages of CO poisoning.⁵¹

The carbon monoxide in waterpipe tobacco smoke results from the combustion of coal, which is why CO levels are high, even when replacing the tobacco with herbal variants. In addition, the burning of coals or 'natural coals' (such as those made from coconuts) also produces significant levels of carcinogenic polycyclic aromatic hydrocarbons (PAHs).⁵²

Chronic exposure to CO may have implications in maintaining blood oxygenation, a concern for surgeons and anaesthetists,⁵³ and is a risk factor in itself for cardiovascular disease.

A final potential health threat, created from the sharing of waterpipes in close proximity between users, is the possibility of infection transmission, in particular tuberculosis,⁵⁴ but other organisms have been shown to exist inside the waterpipe apparatus.⁵⁵

Waterpipe tobacco smoking

A review of the literature concluded that waterpipe smoking can result in levels of up to ten cigarettes' worth of nicotine among people using a waterpipe daily.⁵⁶

Although studies on the health effects of waterpipe tobacco smoking tend to be lacking or of low quality, various detrimental health effects from waterpipe smoking have been established.

A recent systematic review could not rule out associations with:

- bladder cancer;
- nasopharyngeal cancer (cancer of the upper throat);
- oesophageal (gullet) cancer; or
- infertility⁵⁷

High carcinoembryonic antigen levels (CEA) levels, implicated in the spread of cancers, have also been documented amongst those who use waterpipes.⁵⁸

Further evidence shows similar effects of waterpipe tobacco smoking and cigarette smoking on lung function^{59,60} and waterpipe tobacco smoking has also been implicated in the development of chronic obstructive pulmonary disease (COPD).⁶¹

Although there is relatively little evidence in the way of long-term health effects on the heart, acute effects of waterpipe tobacco smoking include increases in heart rate and systolic/diastolic blood pressure^{62–64} as well as impaired baroreflex (the mechanism for regulating blood pressure) sensitivity.⁶⁵ Moreover, 'herbal' waterpipe use can also contribute to such impairment of pulse and blood pressure.⁶⁶

Waterpipe tobacco smoking

Dependence and cessation

As waterpipes can deliver levels of nicotine similar to cigarettes, levels which are considered to lie above the 'addiction threshold',⁶⁷ we can conclude that waterpipe tobacco smoking has the potential to induce dependence.⁵⁶

However, the pattern of dependence is likely to be different from that of cigarettes. It is likely to be shaped by the different mechanisms of delivering nicotine, social context,^{12,68} and sensory cues (aromatic smell, flavoured taste, decorative sight, bubbling sound), as well as different individual smoking behaviour and waterpipe tobacco brand preference⁶⁹ to that of cigarettes.

Furthermore, waterpipe tobacco smoking has been shown to suppress recognised nicotine withdrawal symptoms and cravings in people who smoke, and so has the ability to undermine cigarette smoking cessation attempts.^{70,71}

In response to the growing body of evidence that people who regularly smoke waterpipe tobacco may display or report signs of addiction,^{67,72–75} a validated waterpipe-specific scale to address dependence has been established,⁷⁶ which may be of use to services working with waterpipe users. Among waterpipe café users in London nearly half showed signs of dependence, which was associated with Arab ethnicity, daily waterpipe smoking in the past, and longer waterpipe smoking sessions.⁷⁷ It should be noted however that this scale is likely to need re-validation for appropriate future use in a Western population.

For those who have wanted to stop their waterpipe use, common motivating reasons include health reasons,^{78,68} yet some people who use a waterpipe quit in order to smoke cigarettes instead.^{78,70}

However, the lack of evidence on waterpipe cessation is exemplified by a recent Cochrane review that identified no interventions in the literature and thus the potential use of effective treatments for cigarette dependence, such as nicotine replacement therapy (NRT), remains unproven.⁷⁹ The only evidence for waterpipe cessation can be found from a study in Pakistan which showed that either behavioural support or behavioural support plus seven weeks of bupropion therapy, may be effective cessation methods among people using a waterpipe to smoke tobacco.⁸⁰

Waterpipe tobacco smoking

Advice for people who use a waterpipe to smoke tobacco

Healthcare professionals are in an excellent position to provide information and effective advice, particularly on cessation, for waterpipe users.

In the absence of specific guidelines for the treatment of waterpipe tobacco smoking, and some (limited) evidence of deleterious health effects of waterpipe smoking, it would seem reasonable to adopt the precautionary principle.

Health promotion advocates who engage with the general population or with media should aim to become comfortable and confident when speaking about waterpipe smoking. This is especially true for youth workers and anyone who interacts with young people, given the high and increasing prevalence of waterpipe smoking in this age group. One common pitfall is numerically comparing waterpipe tobacco smoking to cigarette smoking, like-for-like. Several media outlets report that waterpipe tobacco smoking is equivalent to smoking 100 cigarettes or more, however this is not the case and health promoters should avoid potentially confusing numerical comparisons.⁸¹

Asking specifically about waterpipe smoking during consultations where appropriate, will provide an opportunity to advise or refer for cessation support.

A greater desire to quit is seen amongst those given health information compared with those who received no information,⁸² and all health professionals should inform waterpipe users about the potential health risks, and continue to advocate a message of 'no combustible tobacco use in any form' as part of both general health promotion, and in addition to providing tailored support.

It is also recommended that local stop smoking service providers include waterpipe smoking in their treatment protocols.

Waterpipe tobacco smoking

Regulation

The waterpipe industry operates relatively freely and there is considerable scope for better legislation and regulation.

There are strict regulations in the US on prohibiting flavoured cigarettes (except menthol), which has not been extended to other forms of tobacco including the waterpipe.⁸³ Moreover, waterpipe tobacco packaging does not routinely display adequate or relevant health warnings, being non-compliant with the packaging requirements of the World Health Organisation Framework Convention for Tobacco Control (FCTC).⁸⁴

In addition, health warnings are seldom seen on the websites of waterpipe retailers⁸⁵ and their marketing strategies are grossly misleading.⁸⁶ It is worrisome that nicotine labelling on waterpipe Mo'assel tobacco packaging does not correlate with nicotine delivery,⁸⁷ that herbal waterpipe variants deliver as many toxicants to the user as ordinary waterpipe tobacco and that electronic waterpipes have reached the market,⁷ all claiming to reduce harm.

A result of this marketing freedom has been the rise in the number of waterpipe cafes.¹⁵ It is likely that increased accessibility to these venues may encourage initiation,⁸² especially around educational establishments^{16,33} where regular use of these cafes has created suggestions that they serve a community purpose for young people to safely meet, gather, and 'stay out of trouble'.¹⁹

We believe that waterpipe tobacco smoking legislation should, at a minimum, be placed on par with cigarette smoking.

Waterpipe tobacco smoking should be taxed accordingly to discourage purchase, and packaging that is non-compliant with the FCTC should be prohibited. As commercial waterpipe venues do not traditionally display waterpipe tobacco packages to their customers (the pipe is provided pre-packed with tobacco), salient pictorial health warnings should be visible on the apparatus and related accessories.

Waterpipe tobacco smoking

Notes

The main limitation we have faced in drafting this briefing is the lack of reliable and good quality evidence in the literature, especially on the subject of cessation which is the main driver for public health interventions.

In addition, the review is limited by the fact that there is no singular waterpipe product. Many different forms exist each with their own history and method of action, factors which are likely to influence its impact on health, beliefs and attitudes associated with its use, as well possibly requiring specific and different interventions to reduce both use and harm.

Waterpipe tobacco smoking

References

1. Jawad M, McEwen A, McNeill et al. To what extent should waterpipe tobacco smoking become a public health priority? *Addiction* 2013;**108**(11):1873–84
2. Maziak W, Ward KD, Afifi RA, et al. Standardizing questionnaire items for the assessment of waterpipe tobacco use in epidemiological studies. *Public Health* 2005;**119**(5):400–04
3. Maziak W, Ward KD, Afifi RA, et al. Tobacco smoking using a waterpipe: A re-emerging strain in a global epidemic. *Tobacco Control* 2004;**13**(4):327–33
4. Knishkowsky B, Amitai Y. Water-pipe (narghile) smoking: an emerging health risk behavior. *Pediatrics* 2005;**116**(1):e113–e19
5. She J, Yang P, Bai C. Chinese Waterpipe Smoking: A New Risk Factor for Lung Cancer and COPD? *Chest* 2012;**142**(4)
6. Shihadeh A, Salman R, Jaroudi E, et al. Does switching to a tobacco-free waterpipe product reduce toxicant intake? A crossover study comparing CO, NO, PAH, volatile aldehydes, 'tar' and nicotine yields. *Food and Chemical Toxicology* 2012;**50**(5):1494–98
7. Lee YO, Mukherjee A, Grana R. Hookah steam stones: smoking vapour expands from electronic cigarettes to waterpipe. *Tobacco Control* 2012
8. Chaouachi K, Sajid KM. A critique of recent hypotheses on oral (and lung) cancer induced by water pipe (hookah, shisha, narghile) tobacco smoking. *Medical Hypotheses* 2010;**74**(5):843–46
9. Braun R, T G, J W, et al. Hookah Use Among College Students from a Midwest University. *Journal of Community Health* 2011;**37**(2):294–98
10. Fielder RL, Carey KB, Carey MP. Prevalence, frequency, and initiation of hookah tobacco smoking among first-year female college students: A one-year longitudinal study. *Addictive Behaviors* 2012;**37**(2):221–24
11. Ahmed B, Jacob III P, Allen F, et al. Attitudes and practices of hookah smokers in the San Francisco Bay Area. *Journal of Psychoactive Drugs* 2011;**43**(2):146–52
12. Asfar T, Ward KD, Eissenberg T, et al. Comparison of patterns of use, beliefs, and attitudes related to waterpipe between beginning and established smokers. *BMC public health* 2005;**5**
13. Shihadeh A. Investigation of mainstream smoke aerosol of the argileh water pipe. *Food and Chemical Toxicology* 2003;**41**(1):143–52
14. Jawad et al 2013, A qualitative analysis among regular waterpipe tobacco smokers in London universities
INT J TUBERC LUNG DIS 17(10):1364–1369
15. BHF. Rise in 'shisha bars' prompts warning on dangers of waterpipe smoking. Secondary Rise in 'shisha bars' prompts warning on dangers of waterpipe smoking 2012. <http://www.bhf.org.uk/default.aspx?page=14417>.
16. Sutfin EL, McCoy TP, Reboussin BA, et al. Prevalence and correlates of waterpipe tobacco smoking by college students in North Carolina. *Drug and Alcohol Dependence* 2011;**115**(1–2):131–36
17. Nakkash RT, Khalil J, Afifi RA. The rise in narghile (shisha, hookah) waterpipe tobacco smoking: a qualitative study of perceptions of smokers and non smokers. *BMC public health* 2011;**11**
18. Griffiths MA, Harmon TR, Gilly MC. Hubble Bubble Trouble: The Need for Education About and Regulation of Hookah Smoking. *Journal of Public Policy & Marketing* 2011;**30**(1):119–32
19. Schuster RM, Hertel AW, Mermelstein R. Cigar, Cigarillo, and Little Cigar Use Among Current Cigarette-Smoking Adolescents. *Nicotine and Tobacco Research* 2012
20. Sameer uR, Sadiq MA, Parekh MA, et al. Cross-sectional study identifying forms of tobacco used by shisha smokers in Pakistan. *Journal of the Pakistan Medical Association* 2012;**62**(2):192–95
21. Sterling K, Berg CJ, Thomas AN, et al. Factors Associated With Small Cigar Use Among College Students. *American Journal of Health Behavior* 2013;**37**(3):325–33

Waterpipe tobacco smoking

22. Jarrett T, Blosnich J, Tworek C, et al. Hookah Use Among U.S. College Students: Results From the National College Health Assessment II. *Nicotine and Tobacco Research* 2012;**14**(10):1145–53
23. Ward KD, Vander MW, Relyea G, et al. Waterpipe smoking among American military recruits. *Preventive Medicine* 2006;**43**(2):92–97
24. Varsano S, Ganz I, Eldor N, et al. Water-pipe tobacco smoking among school children in Israel: Frequencies, habits, and attitudes. *Harefuah* 2003;**142**(11):736–41+807
25. Jawad M, Abass J, Hariri A, et al. Waterpipe smoking prevalence and attitudes amongst medical students in London. *International Journal of Tuberculosis and Lung Disease* 2013;**17**(1):1–4
26. Cheron-Launay M, Baha M, Mautrait C, et al. Identifying addictive behaviors among adolescents: A school-based survey [French] Reperer les comportements addictifs chez les adolescents: enquete en milieu lyceen. *Archives de Pediatrie* 2011;**18**(7):737–44
27. Jawad M, Khaki H, Hamilton F. Shisha guidance for GPs: Eliciting the hidden history. *British Journal of General Practice* 2012;**62**(595):66–67
28. Akl EA, Gunukula SK, Aleem S, et al. The prevalence of waterpipe tobacco smoking among the general and specific populations: a systematic review. *BMC public health* 2011;**11**
29. El-Rouieheb Z, Tamim H, Kanj M, et al. Cigarette and waterpipe smoking among Lebanese adolescents, a cross-sectional study, 2003–2004. *Nicotine and Tobacco Research* 2008;**10**(2):309–14
30. Al-Lawati JA, Muula AS, Hilmi SA, et al. Prevalence and Determinants of Waterpipe Tobacco Use among Adolescents in Oman. *Sultan Qaboos University Medical Journal* 2008;**8**(1):37–43
31. Mandil A, Hussein A, Omer H, et al. Characteristics and risk factors of tobacco consumption among University of Sharjah students, 2005. *Eastern Mediterranean Health Journal* 2007;**13**(6):1449–58
32. Azab M, Khabour F, Alkaraki K, et al. Water pipe tobacco smoking among university students in Jordan. *Nicotine & Tobacco Research* 2010;**12**(6):606–12
33. Jawad M, Lee JT, Wilson A, et al. Waterpipe and cigarette smoking among secondary school students in London: a comparison of prevalence, beliefs, cessation and predictors. *Nicotine and Tobacco Research*. 2013. [Epub ahead of print]
34. Jackson D, Aveyard P. Waterpipe smoking in students: prevalence, risk factors, symptoms of addiction, and smoke intake. Evidence from one British university. *BMC public health* 2008;**8**
35. Munshi M, Breen R. Waterpipe Smoking amongst London Medical Students. San Diego Conference Centre, 2009.
36. Sajid KM, Akhter M, Malik GQ. Carbon monoxide fractions in cigarette and hookah (hubble bubble) smoke. *JPMA. The Journal of the Pakistan Medical Association* 1993;**43**(9):179–82
37. Saleh R, Shihadeh A. Elevated toxicant yields with narghile waterpipes smoked using a plastic hose. *Food and Chemical Toxicology* 2008;**46**(5):1461–66
38. Shihadeh A, Azar S. A closed-loop control 'playback' smoking machine for generating mainstream smoke aerosols. *Journal of Aerosol Medicine: Deposition, Clearance, and Effects in the Lung* 2006;**19**(2):137–47
39. IARC. Agents Classified by the IARC Monographs, Volumes 1–106. Secondary Agents Classified by the IARC Monographs, Volumes 1–106 2012. <http://monographs.iarc.fr/ENG/Monographs/vol84/mono84-1.pdf>.
40. Al-Kazwini A, Said AJ, Al-Kharouf S. Determination of natural and artificial radionuclide materials in the Ma'assel used in hubbly-bubbly (Shisha) in Jordan. *Toxicological and Environmental Chemistry* 2012;**94**(7):1294–303
41. Khater AEM, Abd NS, Al-Sewaidan HA, et al. Radiological hazards of Narghile (hookah, shisha, goza) smoking: activity concentrations and dose assessment. *Journal of Environmental Radioactivity* 2008;**99**(12):1808–14
42. Radwan G, Hecht SS, Carmella SG, et al. Tobacco-Specific Nitrosamine Exposures in Smokers and Nonsmokers Exposed to Cigarette or Waterpipe Tobacco Smoke. *Nicotine and Tobacco Research* 2012
43. Ashurst JV, Urquhart M, Cook MD. Carbon Monoxide Poisoning Secondary to Hookah Smoking. *The Journal of American Osteopathic Association* 2012;**112**(10):686–88

Waterpipe tobacco smoking

44. Clarke SFJ, Stephens C, Farhan M, et al. Multiple Patients with Carbon Monoxide Toxicity from Water-Pipe Smoking. *Prehospital and Disaster Medicine* 2012;**27**(6):612–14
45. La G, Weiser G, Steiner IP, et al. Carbon monoxide poisoning in narghile (water pipe) tobacco smokers. *CJEM* 2012;**14**(1):57–59
46. Arziman I, Acar YA, Yildirim AO, et al. Five cases of carbon monoxide poisoning due to narghile (shisha). *Hong Kong Journal of Emergency Medicine* 2011;**18**(4):254–57
47. Hojer J, Enghag M. Carbon monoxide poisoning caused by water pipe smoking. *Clinical Toxicology* 2011;**49**(7):702–03
48. Lim BL, Lim GH, Seow E. Case of carbon monoxide poisoning after smoking shisha. *International Journal of Emergency Medicine* 2009;**2**(2):121–22
49. Turkmen S, Eryigit U, Sahin A, et al. Carbon monoxide poisoning associated with water pipe smoking. *Clinical Toxicology* 2011;**49**(7):697–98
50. Cavus UY, Rehber ZH, Ozeke O, et al. Carbon monoxide poisoning associated with Narghile use. *Emergency Medicine Journal* 2010;**27**(5):406
51. Barnett TE, Curbow BA, Soule EK, et al. Carbon monoxide levels among patrons of hookah cafes. *American Journal of Preventive Medicine* 2011;**40**(3):324–28
52. Sepetdjian E, Shihadeh A, Saliba NA. Measurement of 16 polycyclic aromatic hydrocarbons in narghile waterpipe tobacco smoke. *Food and Chemical Toxicology* 2008;**46**(5):1582–90
53. Kesner KL, Ramaiah VK, Hemmer LB, et al. Anesthesia implications of waterpipe use. *Journal of Clinical Anesthesia* 2012;**24**(2):137–40
54. Munckhof WJ, Konstantinos A, Wamsley M, et al. A cluster of tuberculosis associated with use of a marijuana water pipe. *International Journal of Tuberculosis and Lung Disease* 2003;**7**(9):860–65
55. Khoury A, Hanan HS, Hamade H, et al. Characteristics of bacteria in smoking tube of water pipe (Narghile). *European Respiratory Journal*, 2006.
56. Neergaard J, Singh P, Job J, et al. Waterpipe smoking and nicotine exposure: A review of the current evidence. *Nicotine and Tobacco Research* 2007;**9**(10):987–94
57. Akl EA, Gaddam S, Gunukula SK, et al. The effects of waterpipe tobacco smoking on health outcomes: A systematic review. *International Journal of Epidemiology* 2010;**39**(3):834–57
58. Sajid KM, Parveen R, Durr-e-Sabih, et al. Carcinoembryonic antigen (CEA) levels in hookah smokers, cigarette smokers and non-smokers. *JPMA. The Journal of the Pakistan Medical Association* 2007;**57**(12):595–99
59. Raad D, Gaddam S, Schunemann HJ, et al. Effects of water-pipe smoking on lung function: A systematic review and meta-analysis. *Chest* 2011;**139**(4):764–74
60. Boskabady MH, Farhang L, Mahmodinia M, et al. Comparison of pulmonary function and respiratory symptoms in water pipe and cigarette smokers. *Respirology* 2012;**17**(6):950–56
61. Salameh P, Waked M, Khoury F, et al. Waterpipe smoking and dependence are associated with chronic bronchitis: A case-control study in Lebanon [English;French] Narguile et dependance et association avec la bronchite chronique: Etude cas-temoins au Liban. *Eastern Mediterranean Health Journal* 2012;**18**(10):996–1004
62. Al-Safi SA, Ayoub NM, Albalas MA, et al. Does shisha smoking affect blood pressure and heart rate? *Journal of Public Health* 2009;**17**(2):121–26
63. Hakim F, Hellou E, Goldbart A, et al. The acute effects of water-pipe smoking on the cardiorespiratory system. *Chest* 2011;**139**(4):775–81
64. Blank MD, Cobb CO, Kilgalen B, et al. Acute effects of waterpipe tobacco smoking: A double-blind, placebo-control study. *Drug and Alcohol Dependence* 2011;**116**(1–3):102–09
65. Al-Kubati M, Al-Kubati AS, al'Absi M, et al. The short-term effect of water-pipe smoking on the baroreflex control of heart rate in normotensives. *Autonomic Neuroscience: Basic and Clinical* 2006;**126–127**:146–49

Waterpipe tobacco smoking

66. Cobb CO, Sahmarani K, Eissenberg T, et al. Acute toxicant exposure and cardiac autonomic dysfunction from smoking a single narghile waterpipe with tobacco and with a "healthy" tobacco-free alternative. *Toxicology Letters* 2012;**215**(1):70–75
67. Benowitz NL, Henningfield JE. Establishing a Nicotine Threshold for Addiction – The Implications for Tobacco Regulation. *New England Journal of Medicine* 1994;**331**(2):123–25 doi: doi:10.1056/NEJM199407143310212 [published Online First: Epub Date]
68. Ward KD, Hammal F, VanderWeg MW, et al. Are waterpipe users interested in quitting? *Nicotine and Tobacco Research* 2005;**7**(1):149–56
69. Maziak W, Eissenberg T, Ward KD. Patterns of waterpipe use and dependence: Implications for intervention development. *Pharmacology Biochemistry and Behavior* 2005;**80**(1):173–79
70. Al-Dabbagh S, Al-Sinjari KM. Knowledge, attitude and believes of Nargila (hubble-bubble) smoking in Iraq. *Journal of the Bahrain Medical Society* 2005;**17**(2):128–34
71. Rastam S, Eissenberg T, Ibrahim I, et al. Comparative analysis of waterpipe and cigarette suppression of abstinence and craving symptoms. *Addictive Behaviors* 2011;**36**(5):555–59
72. Jawaid A, Zafar AM, Rehman TU, et al. Knowledge, attitudes and practice of university students regarding waterpipe smoking in Pakistan. *International Journal of Tuberculosis and Lung Disease* 2008;**12**(9):1077–84
73. Hammal F, Mock J, Ward KD, et al. A pleasure among friends: how narghile (waterpipe) smoking differs from cigarette smoking in Syria. *Tobacco Control* 2008;**17**(2)
74. Auf RA, Radwan GN, Loffredo CA, et al. Assessment of tobacco dependence in waterpipe smokers in Egypt. *International Journal of Tuberculosis and Lung Disease* 2012;**16**(1):132–37
75. Salameh P, Khayat G, Waked M. Lower Prevalence of Cigarette and Waterpipe Smoking, But a Higher Risk of Waterpipe Dependence in Lebanese Adult Women Than in Men. *Women & Health* 2012;**52**(2):135–50
76. Salameh P, Waked M, Aoun Z. Waterpipe smoking: Construction and validation of the Lebanon Waterpipe Dependence Scale (LWDS-11). *Nicotine and Tobacco Research* 2008;**10**(1):149–58
77. Kassim S, Al-Bakri A, al'Absi M et al. Waterpipe Tobacco Dependence in U.K. Male Adult Residents: A Cross-Sectional Study. *Nicotine and Tobacco Research* 2013
78. Smith JR, Novotny TE, Edland SD, et al. Determinants of Hookah Use among High School Students. *Nicotine and Tobacco Research* 2011;**13**(7):565–72
79. Maziak W, Ward KD, Eissenberg T. Interventions for waterpipe smoking cessation. *Cochrane database of systematic reviews (Online)* 2007(4)
80. Dogar O, Jawad M, Shah SK, Newell JN, Kanaan M, Ahmad M, Khan MA, Siddiqi S. The effect of cessation interventions on hookah smoking: post-hoc analysis of a cluster randomised controlled trial. In press. *Nicotine and Tobacco Research*.
81. Jawad M, Bakir AM, Ali M, Jawad S, Akl EA. Key health themes and reporting of numerical cigarette-waterpipe equivalence in online news articles reporting on waterpipe tobacco smoking: A content analysis. *Tobacco Control*. 2013. [Epub ahead of print]
82. Lipkus IM, Eissenberg T, Schwartz-Bloom RD, et al. Affecting perceptions of harm and addiction among college waterpipe tobacco smokers. *Nicotine and Tobacco Research* 2011;**13**(7):599–610
83. Morris DS, Fiala SC, Pawlak RP. Opportunities for Policy Interventions to Reduce Youth Hookah Smoking in the United States. *Preventing chronic disease* 2012;**9**(120082) doi: <http://dx.doi.org/10.5888/pcd9.120082>[published Online First: Epub Date].
84. Nakkash R, Khalil J. Health warning labelling practices on narghile (shisha, hookah) waterpipe tobacco products and related accessories. *Tobacco Control* 2010;**19**(3):235–39
85. Primack BA, Rice KR, Shensa A, et al. US hookah tobacco smoking establishments advertised on the internet. *American Journal of Preventive Medicine* 2012;**42**(2):150–56
86. Khalil J, Heath RL, Nakkash RT, et al. The tobacco health nexus? Health messages in narghile advertisements. *Tobacco Control* 2009;**18**:420–21
87. Vansickel AR, Shihadeh A, Eissenberg T. Waterpipe tobacco products: Nicotine labelling versus nicotine delivery. *Tobacco Control* 2012;**21**(3):377–79